

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	GRM4
Synonyms	GPRC1D, MGLUR4, mGlu4
Description	Human GRM4 full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q14833
Expression Host	HEK293
Protein Families	GPCR, Transmembrane, Druggable Genome, GPCRDB Class C Metabotropic glutamate pheromone, G-Protein Coupled Receptors Signaling Pathway,
Protein Pathways	
Molecular Weight	The human full length GRM4 protein has a MW of 101.9kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	L-glutamate is the major excitatory neurotransmitter in the central nervous system and activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors, that have been divided into 3 groups on the basis of sequence homology, putative signal transduction mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5 and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3 while Group III includes GRM4, GRM6, GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic AMP cascade but differ in their agonist selectivities. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Feb 2012]
Usage	Research use only
Conjugate	Unconjugated

