

PRODUCT INFORMATION

| | |
|---|---|
| Tag | C-Flag&Strep Tag |
| Expression Host | HEK293 |
| Target | GRM5 |
| Synonyms | GPRC1E, MGLUR5, PPP1R86, mGlu5 |
| Description | Human GRM5-Strep full length protein-synthetic nanodisc |
| Uniprot ID | P41594 |
| Protein Families | GPCR,Transmembrane,Druggable Genome, |
| Protein Pathways | GPCRDB Class C Metabotropic glutamate pheromone, |
| Molecular Weight | The human full length GRM5-Strep protein has a MW of 132.5 kDa |
| Delivery | 6~8weeks |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. |
| Sterility | Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use. |
| Storage&Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene encodes a member of the G-protein coupled receptor 3 protein family. The encoded protein is a metabotropic glutamate receptor, whose signaling activates a phosphatidylinositol-calcium second messenger system. This protein may be involved in the regulation of neural network activity and synaptic plasticity. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. A pseudogene of this gene has been defined on chromosome 11. Alternative splicing results in multiple transcript variants. [provided by RefSeq, May 2014] |
| Usage | Research use only |
| Conjugate | Unconjugated |

