

## PRODUCT INFORMATION

<b>Target</b>	IGF-1R
<b>Synonyms</b>	IGFR, CD221, IGFR, JTK13
<b>Description</b>	Recombinant human IGF-1R Protein with C-terminal human Fc tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P08069
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-Human Fc tag
<b>Molecular Characterization</b>	IGF-1R(Glu31-Asn932) hFc(GLU99-ALA330)
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 129.0 kDa after removal of the signal peptide.
<b>Purity</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Sterility</b>	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
<b>Background</b>	This receptor binds insulin-like growth factor with a high affinity. It has tyrosine kinase activity. The insulin-like growth factor I receptor plays a critical role in transformation events. Cleavage of the precursor generates alpha and beta subunits. It is highly overexpressed in most malignant tissues where it functions as an anti-apoptotic agent by enhancing cell survival. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, May 2014]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated







Figure 1. Human IGF-1R Protein, hFc Tag on SDS-PAGE under reducing condition.

### Human IGF-1R, hFc Tagged Protein ELISA

0.2 µg of Human IGF-1R, hFc tagged protein per well




### Anti-IGF-1R antibody(DMC721), IgG1 Chimeric mAb (ng/mL)

Figure 1. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Human IGF-1R Protein, hFc Tag (PME101889) can bind Anti-IGF-1R antibody(DMC721), IgG1 Chimeric mAb (DMC100721) in a linear range of 8-40 ng/mL.



## Human IGF-1R, hFc Tagged Protein ELISA

0.2  $\mu$ g of Human IGF-1R, hFc tagged protein per well



**Anti-IGF-1R antibody(DMC721), IgG1 Chimeric mAb (ng/mL)**

Figure 2. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human IGF-1R Protein, hFc Tag (PME101889) can bind Anti-IGF-1R antibody(DMC721), IgG1 Chimeric mAb (DMC100721) in a linear range of 8-40 ng/mL.

