

PRODUCT INFORMATION

Target	IL-22RA2
Synonyms	Interleukin-22 Receptor Subunit Alpha-2;IL-22 Receptor Subunit Alpha-2;IL-22R-Alpha-2;IL-22RA2;Cytokine Receptor Class-II Member 10;Cytokine Receptor Family 2 Member 10;CRF2-10;Cytokine Receptor Family Type 2 Soluble 1;CRF2-S1;Interleukin-22-Binding Protein;IL-22BP;IL22BP;ZcytoR16;IL22RA2
Description	Recombinant Human Interleukin-22 Receptor Subunit Alpha-2/Interleukin-22 Binding Protein is produced by our Mammalian expression system and the target gene encoding Thr22-Pro231 is expressed with a Fc tag at the C-terminus.
Delivery	In Stock
Uniprot ID	Q969J5-2
Expression Host	HEK293
Tag	C-Fc Tag
Molecular Characterization	Not available
Molecular Weight	51.8 KDa
Purity	Greater than 95% as determined by reducing SDS-PAGE.
Formulation & Reconstitution	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
Background	Interleukin-22 Receptor Subunit a-2 (IL22RA2) belongs to the type II cytokine receptor family. IL22RA2 is a secreted protein and contains three fibronectin type-III domains. IL22RA2 is widely expressed in many tissues. IL22RA2 functions as an IL22 antagonist and may be important in the regulation of inflammatory response. Three alternatively spliced transcript variants encoding distinct isoforms have been described.
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Greater than 95% as determined by reducing SDS-PAGE.

DIMABIO CONFIDENTIAL

