

## **PRODUCT INFORMATION**

Target IL-37

Interleukin-37;FIL1 Zeta;IL-1X;Interleukin-1 Family Member 7;IL-1F7;Interleukin-1 Homolog 4;IL-1H;IL-1H4;Interleukin-1 Zeta;IL-1 Zeta;Interleukin-1-Related Protein;IL-1RP1;Interleukin-23;IL-37;IL37;FIL1Z;IL1F7;IL1H4;IL1RP1 Recombinant Human Interleukin-37 is produced by our E.coli expression system and the target gene encoding Lys53-Asp218 is expressed Svnonvms

expressed.

**Delivery** In Stock O9NZH6 **Uniprot ID Expression Host** E.coli

Tag

**Description** 

Molecular Not available Characterization **Molecular Weight** 18.7 KDa

**Purity** Greater than 95% as determined by reducing SDS-PAGE.

Lyophilized from a 0.2  $\mu m$  filtered solution of 20mM PB, 150mM NaCl, 2mM DTT, pH 7.4. Formulation &

Reconstitution

Store at -20°C to -80°C for 12 months in lyophilized form. After Storage & Shipping

reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized

proteins are shipped at ambient temperature.

Human Interleukin family 1 Member 7 (IL1F7) is a member of the Interleukin 1 cytokine family. Five alternatively spliced transcript variants encoding distinct isoforms have been reported with distinct expression profiles. The longest IL1F7 transcript, referred to as IL1F7b or IL1F7 isoform 1, encodes a 218 amino acid residues proprotein containing a 45 amino acid propeptide, which is cleaved to generate mature protein. IL1F7b binds to IL18 Ra with low affinity but does not even any IL18 agonities or

**Background** 

with low affinity but does not exert any IL18 agonistic or antagonistic effects. IL1F7b also binds interleukin 18 binding protein (IL-18BP), an inhibitory binding protein of interleukin 18 (IL-18), and subsequently forms a complex with IL18 receptor beta subunit, and through which it inhibits the activity of IL-18.

Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)








Figure 1. Greater than 95% as determined by reducing SDS-PAGE.



Email: info@dimabio.com Website: www.dimabio.com