

## **PRODUCT INFORMATION**

**IL13 Target** 

**Synonyms** IL-13;NC30

Recombinant human IL13 protein with C-terminal **Description** 

human Fc tag

**Delivery** In Stock **Uniprot ID** P35225 **Expression Host HEK293** 

Tag C-Human Fc Tag

Molecular

Purity

**Background** 

IL13(Thr21-Leu132) hFc(Glu99-Ala330) Characterization

The protein has a predicted molecular mass of **Molecular Weight** 

38.1 kDa after removal of the signal peptide. The apparent molecular mass of IL13-hFc is

approximately 35-70 kDa due to glycosylation.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & Reconstitution

- 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes an immunoregulatory cytokine produced primarily by activated Th2 cells. This cytokine is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine down-regulates macrophage activity, thereby inhibits the production of pro-inflammatory cytokines and chemokines. This cytokine is found

to be critical to the pathogenesis of allergen-induced asthma but operates through mechanisms independent of IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine particularly close to IL4. [provided by RefSeq, Jul 2008]

> Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only

Conjugate Unconjugated








Figure 1. Human IL13 Protein, hFc Tag on SDS-PAGE under reducing condition.



