

PRODUCT INFORMATION

Target	IL17C
Synonyms	CX2;IL-17C
Description	Recombinant Human IL17C with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	Q9P0M4
Expression Host	HEK293
Tag	C-Human Fc Tag
Molecular Characterization	IL17C(His19-Val197) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 45.9 kDa after removal of the signal peptide. The apparent molecular mass of IL17C-hFc is approximately 35-55 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
Background	The protein encoded by this gene is a T cell-derived cytokine that shares the sequence similarity with IL17. This cytokine was reported to stimulate the release of tumor necrosis factor alpha and interleukin 1 beta from a monocytic cell line. The expression of this cytokine was found to be restricted to activated T cells. [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human IL17C Protein, hFc Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

