

PRODUCT INFORMATION

Target	IL20RA
Synonyms	CRF2-8, IL-20R1, IL-20RA, IL-20R-alpha
Description	Recombinant human IL20RA Protein with C-terminal 10×His tag
Delivery	In Stock
Uniprot ID	Q9UHF4
Expression Host	HEK293
Tag	C-10×His tag
Molecular Characterization	IL20RA(Val30-Lys250) 10×His tag
Molecular Weight	The protein has a predicted molecular mass of 26.6 kDa after removal of the signal peptide.
Purity	The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a member of the type II cytokine receptor family. The encoded protein is a subunit of the receptor for interleukin 20, a cytokine that may be involved in epidermal function. The interleukin 20 receptor is a heterodimeric complex consisting of the encoded protein and interleukin 20 receptor beta. This gene and interleukin 20 receptor beta are highly expressed in skin, and are upregulated in psoriasis. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2013]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human IL20RA Protein, His Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

