

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	ILDR2
Synonyms	C1orf32; dJ782G3.1
Description	Human ILDR2 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	Q71H61
Expression Host	HEK293
Protein Families	Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length ILDR2 protein has a MW of 71.2 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Background	May be involved in ER stress pathways with effects on lipid homeostasis and insulin secretion. With ILDR1 and LSR, involved in the maintain of the epithelial barrier function through the recruitment of MARVELD2/tricellulin to tricellular tight junctions. Also functions as a B7-like protein family member expressed on immune cells and inflamed tissue and with T-cell inhibitory activity.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C(Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate ILDR2-Nanodisc
0.2 μ g Human ILDR2-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag ILDR2-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with ILDR2-Nanodisc is 4.527ng/ml.



Figure2. WB analysis of Human ILDR2-Nanodisc with anti-Flag monoclonal antibody at 1/5000 dilution, followed by Goat Anti-Rabbit IgG HRP at 1/5000 dilution

