

**PRODUCT INFORMATION**

<b>Tag</b>	C-Flag&Strep Tag
<b>Target</b>	KCNH2
<b>Synonyms</b>	ERG-1; ERG1; H-ERG; HERG; HERG1; Kv11.1; LQT2; SQT1
<b>Description</b>	Human KCNH2-Strep full length protein-synthetic nanodisc
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q12809
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Druggable Genome, Ion Channels: Potassium, Transcription Factors, Transmembrane
<b>Protein Pathways</b>	N/A
<b>Molecular Weight</b>	The human full length KCNH2-Strep protein has a MW of 126.7 kDa
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	A voltage-activated potassium channel belonging to the eag family. It shares sequence similarity with the Drosophila ether-a-go-go (eag) gene. Mutations in this gene can cause long QT syndrome type 2 (LQT2).
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated



### ELISA assay to evaluate KCNH2-Strep-Nanodisc 0.2 $\mu$ g Human KCNH2-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag KCNH2-Strep-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with KCNH2-Strep-nanodisc is 2.328ng/ml.

kDa M R

250  
130  
100  
70  
55  
35  
25  
15  
10



Figure 2. Human KCNH2-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

