

PRODUCT INFORMATION

Tag	C-Flag Tag
Expression Host	HEK293
Target	LPAR2
Synonyms	EDG-4, EDG4, LPA-2, LPA2
Description	Human LPAR2 full length protein-synthetic nanodisc
Uniprot ID	Q9HBW0
Protein Families	GPCR, Transmembrane, Druggable Genome,
Protein Pathways	Cancer, G-Protein Coupled Receptors Signaling Pathway,
Molecular Weight	The human full length LPAR2 protein has a MW of 38.7kDa
Delivery	6~8 weeks
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a member of family I of the G protein-coupled receptors, as well as the EDG family of proteins. This protein functions as a lysophosphatidic acid (LPA) receptor and contributes to Ca ²⁺ mobilization, a critical cellular response to LPA in cells, through association with Gi and Gq proteins. An alternative splice variant has been described but its full length sequence has not been determined. [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated

