

PRODUCT INFORMATION

MUC1 Target

ADMCKD;ADMCKD1;CA ADMCKD;ADMCKD1;CA 15-3;CD227;EMA;H23AG;KL-6;MAM6;MCD;MCKD;MCKD1;MUC-1;MUC-1/SEC;MUC-1/X;MUC1/ZD;PEM;PEMT;PUM Synonyms

Description Recombinant Human MUC1 Protein with C-terminal human Fc tag

Delivery Uniprot ID P15941 **Expression Host** HFK293 C-Human Fc Tag

MUC1(Leu1036-Val1156) hFc(Glu99-Ala330)

Molecular Characterization **Molecular Weight** The protein has a predicted molecular mass of 39.7 kDa after removal of the signal peptide.

Purity The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Storage & Shipping

Nophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

This gene encodes a membrane-bound protein that is a member of the mucin family. Mucins are O-glycosylated proteins that play an essential role in forming protective mucous barriers on epithelial surfaces. These proteins also play a role in intracellular signaling. This protein is expressed on the apical surface of epithelial cells that line the mucosal surfaces of many different tissues including lung, breast stomach and pancreas. This protein is protein is proteolytically cleaved into alpha and beta subunits that form a heterodimeric complex. The N-terminal alpha subunit functions in cell-adhesion and the C-terminal beta subunit is involved in cell signaling. Overexpression, aberrant intracellular localization, and changes in glycosylation of this protein have been associated with carcinomas. This gene is known to contain a highly polymorphic variable number tandem repeats (VNTR) domain. Alternate splicing results in multiple transcript variants.

Usage Unconjugated Conjugate

Background




Figure 1. Human MUC1 Protein, hFc Tag on SDS-PAGE under reducing condition.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

