

PRODUCT INFORMATION

Target	NINJ1
Synonyms	NIN1;NINJURIN
Description	Recombinant Human NINJ1 Protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	Q92982
Expression Host	HEK293
Tag	C-Human Fc Tag
Molecular Characterization	NINJ1(Met1-Leu80) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 34.6 kDa after removal of the signal peptide. The apparent molecular mass of NINJ1-hFc is approximately 35-55 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The ninjurin protein is upregulated after nerve injury both in dorsal root ganglion neurons and in Schwann cells (Araki and Milbrandt, 1996 [PubMed 8780658]). It demonstrates properties of a homophilic adhesion molecule and promotes neurite outgrowth from primary cultured dorsal root ganglion neurons.[supplied by OMIM, Aug 2009]
Usage	Research use only






Figure 1.Human NINJ1 Protein, hFc Tag on SDS-PAGE under reducing condition.

