

## PRODUCT INFORMATION

<b>Target</b>	NKG2A
<b>Synonyms</b>	CD159A;NKG2;NKG2A
<b>Description</b>	Recombinant Human NKG2A Protein with N-terminal human Fc tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P26715
<b>Expression Host</b>	HEK293
<b>Tag</b>	N-Human Fc Tag
<b>Molecular Characterization</b>	hFc(Glu99-Ala330) NKG2A(Pro94-Leu233)
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 42.2 kDa after removal of the signal peptide. The apparent molecular mass of hFc-NKG2A is approximately 35-55 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Natural killer (NK) cells are lymphocytes that can mediate lysis of certain tumor cells and virus-infected cells without previous activation. They can also regulate specific humoral and cell-mediated immunity. The protein encoded by this gene belongs to the killer cell lectin-like receptor family, also called NKG2 family, which is a group of transmembrane proteins preferentially expressed in NK cells. This family of proteins is characterized by the type II membrane orientation and the presence of a C-type lectin domain. This protein forms a complex with another family member, KLRD1/CD94, and has been implicated in the recognition of the MHC class I HLA-E molecules in NK cells. The genes of NKG2 family members form a killer cell lectin-like receptor gene cluster on chromosome 12. Multiple alternatively spliced transcript variants encoding distinct isoforms have been observed. [provided by RefSeq, Jan 2015]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated







Figure 1. Human NKG2A Protein, hFc Tag on SDS-PAGE under reducing condition.

### Human NKG2A, hFc Tagged protein ELISA

0.2  $\mu$ g of Human NKG2A, hFc tagged protein per well

Figure 2. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human NKG2A Protein, hFc Tag (PME100851) can bind Anti-NKG2A antibody(DMC298), IgG1 Chimeric mAb in a linear range of 3.20-80 ng/mL.