

PRODUCT INFORMATION

Target	NPR1
Synonyms	ANP-A;ANPR-A;ANPRA;NPR-A;GC-A
Description	Recombinant human NPR1 protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	P16066
Expression Host	HEK293
Tag	C-Human Fc Tag
Molecular Characterization	NPR1(Gly33-Glu473) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 75.0 kDa after removal of the signal peptide. The apparent molecular mass of NPR1-hFc is approximately 95-130 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
Background	Guanylyl cyclases, catalyzing the production of cGMP from GTP, are classified as soluble and membrane forms (Garbers and Lowe, 1994 [PubMed 7982997]). The membrane guanylyl cyclases, often termed guanylyl cyclases A through F, form a family of cell-surface receptors with a similar topographic structure: an extracellular ligand-binding domain, a single membrane-spanning domain, and an intracellular region that contains a protein kinase-like domain and a cyclase catalytic domain. GC-A and GC-B function as receptors for natriuretic peptides; they are also referred to as atrial natriuretic peptide receptor A (NPR1) and type B (NPR2; MIM 108961). Also see NPR3 (MIM 108962), which encodes a protein with only the ligand-binding transmembrane and 37-amino acid cytoplasmic domains. NPR1 is a membrane-bound guanylate cyclase that serves as the receptor for both atrial and brain natriuretic peptides (ANP (MIM 108780) and BNP (MIM 600295), respectively).[supplied by OMIM, May 2009]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human NPR1 Protein, hFc Tag on SDS-PAGE under reducing condition.

