

## **PRODUCT INFORMATION**

C-Flag&Strep Tag Tag

NPY2R **Target Synonyms** NPY2-R

Human NPY2R-Strep full length protein-synthetic **Description** 

nanodisc **Delivery** In Stock **Uniprot ID** P49146 **Expression Host HEK293** 

**Protein Families** Druggable Genome, GPCR, Transmembrane

**Protein Pathways** Neuroactive ligand-receptor interaction

The human full length NPY2R-Strep protein has a **Molecular Weight** 

MW of 42.7 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before Formulation & Reconstitution lyophilization. Please see Certificate of Analysis

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Storage & Shipping

Lyophilized proteins are shipped at ambient

temperature.

Receptor for neuropeptide Y and peptide YY. The

rank order of affinity of this receptor for pancreatic polypeptides is PYY > NPY > PYY (3-36) > NPY (2-36) > [Ile-31, Gln-34] PP > [Leu-31, Pro-34] NPY > PP, [Pro-34] PYY and NPY free acid

Email: info@dimabio.com Website: www.dimabio.com

free acid.

**Usage** Research use only Conjugate Unconjugated

**Background** 





## ELISA assay to evaluate NPY2R-Strep-Nanodisc 0.2µg Human NPY2R-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag NPY2R-Strep-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with NPY2R-Strep-nanodisc is 5.980ng/ml.




Figure 2. Human NPY2R-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

Email: info@dimabio.com Website: www.dimabio.com