

PRODUCT INFORMATION

Target	NRG4
Synonyms	HRG4
Description	Recombinant human NRG4 Protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	Q8WWG1
Expression Host	HEK293
Tag	C-Human Fc tag
Molecular Characterization	NRG4(Met1-Phe62) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 32.9 kDa after removal of the signal peptide.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
Background	The neuregulins, including NRG4, activate type-1 growth factor receptors (see EGFR; MIM 131550) to initiating cell-to-cell signaling through tyrosine phosphorylation (Harari et al., 1999 [PubMed 10348342]).[supplied by OMIM, Mar 2008]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human NRG4 Protein, hFc Tag on SDS-PAGE under reducing condition.

