

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	NTSR1
Synonyms	NTR
Description	Human NTSR1 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	P30989
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Calcium signaling pathway, Neuroactive ligand-receptor interaction
Molecular Weight	The human full length NTSR1 protein has a MW of 46.1 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Formulation & Reconstitution	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Storage&Shipping	Neurotensin receptor 1 belongs to the large superfamily of G-protein coupled receptors. NTSR1 mediates the multiple functions of neurotensin, such as hypotension, hyperglycemia, hypothermia, antinociception, and regulation of intestinal motility and secretion.
Background	
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate NTSR1-Nanodisc
0.2 μ g Human NTSR1-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag NTSR1-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with NTSR1-Nanodisc is 2.583ng/ml.



Figure2. Human NTSR1-Nanodisc, Flag Tag on SDS-PAGE

