

PRODUCT INFORMATION

Target	OPG
Synonyms	Tumor necrosis factor receptor superfamily member 11B;Osteoclastogenesis inhibitory factor;Osteoprotegerin;TNFRSF11B;OCIF;OPG
Description	Recombinant Human Osteoprotegerin is produced by our Mammalian expression system and the target gene encoding Glu22-Leu201 is expressed with a Fc tag at the C-terminus.
Delivery	In Stock
Uniprot ID	O00300
Expression Host	HEK293
Tag	C-Fc Tag
Molecular Characterization	Not available
Molecular Weight	47.2 KDa
Purity	Greater than 95% as determined by reducing SDS-PAGE.
Formulation & Reconstitution	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	TNFRSF11B is a secreted protein, containing 2 death domains and 4 TNFR-Cys repeats. TNFRSF11B is a decoy receptor for the receptor activator of nuclear factor kappa B ligand (RANKL). By binding RANKL, TNFRSF11B inhibits nuclear kappa B (NF-κB) which is a central and rapid acting transcription factor for immune-related genes, and a key regulator of inflammation, innate immunity, and cell survival and differentiation. TNFRSF11B levels are influenced by voltage-dependent calcium channelsCav1.2. TNFRSF11B can reduce the production of osteoclasts by inhibiting the differentiation of osteoclast precursors into osteoclasts and also regulates the resorption of osteoclasts in vitroand in vivo. TNFRSF11B binding to RANKL on osteoblast/stromal cells, blocks the RANKL-RANK ligand interaction between osteoblast/stromal cells and osteoclast precursors. This has the effect of inhibiting the differentiation of the osteoclast precursor into a mature osteoclast.
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Greater than 95% as determined by reducing SDS-PAGE.

DIMABIO CONFIDENTIAL

