

**PRODUCT INFORMATION**

<b>Tag</b>	C-Flag Tag
<b>Expression Host</b>	HEK293
<b>Target</b>	OPRX
<b>Synonyms</b>	KOR-3, KOR3, NOCIR, NOP, NOPr, OOR, OPRL, ORL1, PNOCR
<b>Description</b>	Human OPRX full length protein-synthetic nanodisc
<b>Uniprot ID</b>	P41146
<b>Protein Families</b>	GPCR, Transmembrane, Druggable Genome,
<b>Protein Pathways</b>	GPCRDB Class A Rhodopsin-like, Peptide GPCRs,
<b>Molecular Weight</b>	The human full length OPRX protein has a MW of 40.7kDa
<b>Delivery</b>	6~8weeks
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution
<b>Sterility</b>	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein encoded by this gene is a member of the 7 transmembrane-spanning G protein-coupled receptor family, and functions as a receptor for the endogenous, opioid-related neuropeptide, nociceptin/orphanin FQ. This receptor-ligand system modulates a variety of biological functions and neurobehavior, including stress responses and anxiety behavior, learning and memory, locomotor activity, and inflammatory and immune responses. A promoter region between this gene and the 5'-adjacent RGS19 (regulator of G-protein signaling 19) gene on the opposite strand functions bi-directionally as a core-promoter for both genes, suggesting co-operative transcriptional regulation of these two functionally related genes. Alternatively spliced transcript variants have been described for this gene. A recent study provided evidence for translational readthrough in this gene, and expression of an additional C-terminally extended isoform via the use of an alternative in-frame translation termination codon. [provided by RefSeq, Dec 2017]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated

