

PRODUCT INFORMATION

OR2H1 **Target**

6M1-16; dJ994E9.4; HS6M1-16;

OLFR42A-9004-14; OLFR42A-9004.14/9026.2; **Synonyms**

OR2H6; OR2H8; OR6-2

Human OR2H1 full length protein-synthetic **Description**

nanodisc

Delivery In Stock **Uniprot ID** Q9GZK4 **Expression Host HEK293**

Protein Families Druggable Genome, Transmembrane

Protein Pathways Olfactory transduction

The human full length OR2H1 Protein has a MW of **Molecular Weight**

35.3 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% 8% trehalose is added as protectants before Formulation & Reconstitution lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments. Store at -20°C to -80°C for 12 months in

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and

hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor

genes and proteins for this organism is independent of other organisms.

Usage Research use only

Background

Email: info@dimabio.com Website: www.dimabio.com





ELISA assay to evaluate OR2H1-Nanodisc 0.2μg Human OR2H1-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag OR2H1-Nanodisc ($0.2\mu g/per$ well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with OR2H1-Nanodisc is 1.37ng/ml.




Figure 2. Human OR2H1-Nanodisc, Flag Tag on SDS-PAGE

Email: info@dimabio.com Website: www.dimabio.com