

**PRODUCT INFORMATION**

|   |   |
|---|---|
| <b>Tag</b>                              | C-Flag&Strep Tag  |
| <b>Target</b>                           | OXYR  |
| <b>Synonyms</b>                         | OT-R; OTR   |
| <b>Description</b>                      | Human OXYR-Strep full length protein-synthetic nanodisc   |
| <b>Delivery</b>                         | 6~8weeks  |
| <b>Uniprot ID</b>                       | P30559  |
| <b>Expression Host</b>                  | HEK293  |
| <b>Protein Families</b>                 | Druggable Genome, GPCR, Transmembrane   |
| <b>Protein Pathways</b>                 | Calcium signaling pathway, Neuroactive ligand-receptor interaction  |
| <b>Molecular Weight</b>                 | The human full length OXYR-Strep protein has a MW of 42.8 kDa<br>Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. |
| <b>Formulation &amp; Reconstitution</b> | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| <b>Storage&amp;Shipping</b>             |   |
| <b>Background</b>                       | The protein belongs to the G-protein coupled receptor family and acts as a receptor for oxytocin. Its activity is mediated by G proteins which activate a phosphatidylinositol-calcium second messenger system. The oxytocin-oxytocin receptor system plays an important role in the uterus during parturition.   |
| <b>Usage</b>                            | Research use only   |
| <b>Conjugate</b>                        | Unconjugated  |

