

**PRODUCT INFORMATION**

<b>Tag</b>	C-Flag&Strep Tag
<b>Expression Host</b>	HEK293
<b>Target</b>	P2RX3
<b>Synonyms</b>	P2X3
<b>Description</b>	Human P2RX3-Strep full length protein-synthetic nanodisc
<b>Uniprot ID</b>	P56373
<b>Protein Families</b>	Druggable Genome, Ion Channels: ATP Receptors, Transmembrane
<b>Protein Pathways</b>	Calcium signaling pathway, Neuroactive ligand-receptor interaction
<b>Molecular Weight</b>	The human full length P2RX3-Strep protein has a MW of 44.3 kDa
<b>Delivery</b>	In Stock
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
<b>Sterility</b>	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	A member of the P2X purinergic receptor (purinoceptor) gene family which includes seven members (P2RX1 - P2RX7). P2X purinoceptors are a family of cation-permeable, ligand-gated ion channels that open in response to the binding of extracellular adenosine 5'-triphosphate (ATP). The encoded protein is a subunit of the trimeric P2X3 receptor ion channel which is expressed by sensory or autonomic neurons. A deficiency of the orthologous protein in mice is associated with reduced pain-related behavior and urinary bladder hyporeflexia.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated



### ELISA assay to evaluate P2RX3-Strep-Nanodisc 0.2 $\mu$ g Human P2RX3-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag P2RX3-Strep-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with P2RX3-Strep-nanodisc is 8.046ng/ml.




Figure 2. Human P2RX3-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

