

PRODUCT INFORMATION

PD-1 **Target**

PD1; PDCD1; CD279; SLEB2; hPD-1; hPD-I; hSLE1; **Synonyms**

AIMTBS

Recombinant human PD-1 Protein with C-terminal **Description**

human Fc tag

Delivery In Stock **Uniprot ID** Q15116 **Expression Host HEK293**

Tag C-Human Fc tag

Molecular

Background

PD-1(Leu25-Thr145) hFc(Glu99-Ala330) Characterization

The protein has a predicted molecular mass of **Molecular Weight** 39.7 kDa after removal of the signal peptide.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue **Purity**

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & - 8% trehalose is added as protectants before Reconstitution lyophilization. Please see Certificate of Analysis

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

Programmed cell death protein 1 (PDCD1) is an immune-inhibitory receptor expressed in activated T cells; it is involved in the regulation of T-cell functions, including those of effector CD8+

T cells. In addition, this protein can also promote the differentiation of CD4+ T cells into T regulatory cells. PDCD1 is expressed in many types of tumor to play a relation of the contract of the contract

demonstrated to play a role in anti-tumor immunity. Moreover, this protein has been shown to be involved in safeguarding against


autoimmunity, however, it can also contribute to the inhibition of effective anti-tumor and anti-

> Email: info@dimabio.com Website: www.dimabio.com

microbial immunity. [provided by RefSeq, Aug

Research use only Usage

Conjugate Unconjugated



/+86-400-006-0995(China)








Figure 1. Human PD-1 Protein, hFc Tag on SDS-PAGE under reducing condition.

Human PD1(25-145), hFc Tagged protein ELISA

0.2 μg of Human PD1(25-145), hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human PD1(25-145) Protein, hFc Tag (PME101789) can bind Anti-PD-1(rosnilimab biosimilar) mAb (BME100657) in a linear range of 3.2–8 ng/mL.

Email: info@dimabio.com Website: www.dimabio.com

