

## PRODUCT INFORMATION

<b>Target</b>	PDGFRA
<b>Synonyms</b>	CD140A; PDGFR2; PDGFR-2
<b>Description</b>	Recombinant human PDGFRA Protein with C-terminal 6×His tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P16234
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-6×His tag
<b>Molecular Characterization</b>	PDGFRA(Gln24-Ala528) 6×His tag
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 57.4 kDa after removal of the signal peptide. The apparent molecular mass of PDGFRA-His is approximately 100-130 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a cell surface tyrosine kinase receptor for members of the platelet-derived growth factor family. These growth factors are mitogens for cells of mesenchymal origin. The identity of the growth factor bound to a receptor monomer determines whether the functional receptor is a homodimer or a heterodimer, composed of both platelet-derived growth factor receptor alpha and beta polypeptides. Studies suggest that this gene plays a role in organ development, wound healing, and tumor progression. Mutations in this gene have been associated with idiopathic hypereosinophilic syndrome, somatic and familial gastrointestinal stromal tumors, and a variety of other cancers. [provided by RefSeq, Mar 2012]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated






Figure 1. Human PDGFRA Protein, His Tag on SDS-PAGE under reducing condition.

### Human PDGFRA, His Tagged protein ELISA

0.2 µg of Human PDGFRA, His tagged protein per well




Figure 2. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Human PDGFRA Protein, His Tag (PME101358) can bind Anti-PDGFR(olaratumab biosimilar) mAb (BME100598) in a linear range of 2.29-61.73 ng/mL.

