

PRODUCT INFORMATION

| | |
|---|--|
| Tag | C-Flag Tag |
| Target | PITC1 |
| Synonyms | M-RDGB-beta, MRDGBbeta, RDGB-BETA, RDGBB, RDGBB1 |
| Description | Human PITC1 full length protein-synthetic nanodisc |
| Delivery | 6~8weeks |
| Uniprot ID | Q9UKF7 |
| Expression Host | HEK293 |
| Protein Families | Ion Channels: Other |
| Protein Pathways | N/A |
| Molecular Weight | The human full length PITC1 protein has a MW of 38.4kDa |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution |
| Storage&Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene encodes a member of the phosphatidylinositol transfer protein family. The encoded cytoplasmic protein plays a role in multiple processes including cell signaling and lipid metabolism by facilitating the transfer of phosphatidylinositol between membrane compartments. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene, and a pseudogene of this gene is located on the long arm of chromosome 1. [provided by RefSeq, May 2012] |
| Usage | Research use only |
| Conjugate | Unconjugated |

