

## PRODUCT INFORMATION

|   |   |
|---|---|
| <b>Target</b>                           | PLA2R1  |
| <b>Synonyms</b>                         | CLEC13C;PLA2-R;PLA2G1R;PLA2IR;PLA2R   |
| <b>Description</b>                      | Recombinant Human PLA2R1(21-164 223-359 1097-1246) Protein with C-terminal 10×His tag   |
| <b>Delivery</b>                         | In Stock  |
| <b>Uniprot ID</b>                       | Q13018  |
| <b>Expression Host</b>                  | HEK293  |
| <b>Tag</b>                              | C-10×His Tag  |
| <b>Molecular Characterization</b>       | PLA2R1(Glu21-Lys164 Thr223-Asn359 Glu1097-Leu1246) 10×His tag   |
| <b>Molecular Weight</b>                 | The protein has a predicted molecular mass of 50.3 kDa after removal of the signal peptide. The apparent molecular mass of PLA2R1(21-164 223-359 1097-1246)-10×His is approximately 35-70 kDa due to glycosylation.   |
| <b>Purity</b>                           | The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining.  |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.  |
| <b>Storage&amp;Shipping</b>             | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| <b>Sterility</b>                        | Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.   |
| <b>Background</b>                       | This gene represents a phospholipase A2 receptor. The encoded protein likely exists as both a transmembrane form and a soluble form. The transmembrane receptor may play a role in clearance of phospholipase A2, thereby inhibiting its action. Polymorphisms at this locus have been associated with susceptibility to idiopathic membranous nephropathy. Alternatively spliced transcript variants encoding different isoforms have been identified.[provided by RefSeq, Sep 2010] |
| <b>Usage</b>                            | Research use only   |
| <b>Conjugate</b>                        | Unconjugated  |






Figure 1. Human PLA2R1(21-164 223-359 1097-1246) Protein, His Tag on SDS-PAGE under reducing condition.

