

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	SCN8A
Synonyms	BFIS5, CERIII, CIAT, DEE13, EIEE13, MED, MYOCL2, NaCh6, Nav1.6, PN4
Description	Human SCN8A full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q9UQD0
Expression Host	HEK293
Protein Families	Ion Channels: Sodium
Protein Pathways	N/A
Molecular Weight	The human full length SCN8A protein has a MW of 225.3kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a member of the sodium channel alpha subunit gene family. The encoded protein forms the ion pore region of the voltage-gated sodium channel. This protein is essential for the rapid membrane depolarization that occurs during the formation of the action potential in excitable neurons. Mutations in this gene are associated with cognitive disability, pancerebellar atrophy and ataxia. Alternate splicing results in multiple transcript variants.[provided by RefSeq, May 2010]
Usage	Research use only
Conjugate	Unconjugated

