

PRODUCT INFORMATION

C-Flag&Strep Tag Tag

SLC2A4 **Target** GLUT4 **Synonyms**

Formulation & Reconstitution

Background

Human SLC2A4-Strep full length protein-synthetic **Description**

nanodisc **Delivery** 6~8weeks **Uniprot ID** P14672 **Expression Host HEK293**

Protein Families Druggable Genome, Transmembrane

Adipocytokine signaling pathway, Insulin signaling pathway, Type II diabetes mellitus **Protein Pathways**

The human full length SLC2A4-Strep protein has a **Molecular Weight**

MW of 54.6 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

This gene is a member of the solute carrier family 2 (facilitated glucose transporter) family and encodes a protein that functions as an insulin-regulated facilitative glucose transporter. In the absence of insulin, this integral membrane protein is sequestered within the cells of muscle and adipose tissue. Within minutes of insulin stimulation, the protein moves to the cell surface

and begins to transport glucose across the cell membrane. Mutations in this gene have been associated with noninsulin-dependent diabetes mellitus (NIDDM). [provided by RefSeq, Jul 2008]

> Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only

Conjugate Unconjugated

