

PRODUCT INFORMATION

Target	SPA17
Synonyms	CT22;SP17;SP17-1
Description	Recombinant Human SPA17 Protein with C-terminal 6×His tag
Delivery	In Stock
Uniprot ID	Q15506
Expression Host	HEK293
Tag	C-6×His Tag
Molecular Characterization	SPA17(Met1-Lys151) 6×His tag
Molecular Weight	The protein has a predicted molecular mass of 18.2 kDa after removal of the signal peptide. The apparent molecular mass of SPA17-His is approximately 15-25 kDa due to glycosylation.
Purity	The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a protein present at the cell surface. The N-terminus has sequence similarity to human cAMP-dependent protein kinase A (PKA) type II alpha regulatory subunit (RIIa) while the C-terminus has an IQ calmodulin-binding motif. The central portion of the protein has carbohydrate binding motifs and likely functions in cell-cell adhesion. The protein was initially characterized by its involvement in the binding of sperm to the zona pellucida of the oocyte. Recent studies indicate that it is also involved in additional cell-cell adhesion functions such as immune cell migration and metastasis. A retrotransposed pseudogene is present on chromosome 10q22.[provided by RefSeq, Jan 2009]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human SPA17 Protein, His Tag on SDS-PAGE under reducing condition.

