

**PRODUCT INFORMATION**

|   |   |
|---|---|
| <b>Tag</b>                              | C-Flag&Strep Tag  |
| <b>Target</b>                           | SSR5  |
| <b>Synonyms</b>                         | SS-5-R  |
| <b>Description</b>                      | Human SSR5-Strep full length protein-synthetic nanodisc   |
| <b>Delivery</b>                         | 6~8weeks  |
| <b>Uniprot ID</b>                       | P35346  |
| <b>Expression Host</b>                  | HEK293  |
| <b>Protein Families</b>                 | GPCR,Transmembrane,Druggable Genome,  |
| <b>Protein Pathways</b>                 | GPCRDB Class A Rhodopsin-like,Peptide GPCRs,Metabolic and Obesity,  |
| <b>Molecular Weight</b>                 | The human full length SSR5-Strep protein has a MW of 39.2 kDa<br>Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.   |
| <b>Formulation &amp; Reconstitution</b> | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| <b>Storage&amp;Shipping</b>             | Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.   |
| <b>Sterility</b>                        | Somatostatin and its related peptide cortistatin exert multiple biological actions on normal and tumoral tissue targets by interacting with somatostatin receptors (SSTRs). The protein encoded by this gene is one of the SSTRs, which is a multi-pass membrane protein and belongs to the G-protein coupled receptor 1 family. The activity of this receptor is mediated by G proteins which inhibit adenylyl cyclase, and different regions of this receptor molecule are required for the activation of different signaling pathways. A mutation in this gene results in somatostatin analog resistance. Alternatively spliced transcript variants have been identified in this gene.[provided by RefSeq, Feb 2010] |
| <b>Background</b>                       |   |
| <b>Usage</b>                            | Research use only   |
| <b>Conjugate</b>                        | Unconjugated  |

