

**PRODUCT INFORMATION**

<b>Tag</b>	C-Flag Tag
<b>Expression Host</b>	HEK293
<b>Target</b>	SSTR2
<b>Synonyms</b>	SS-2-R; SS2-R; SS2R; SST2
<b>Description</b>	Human SSTR2 full length protein membrane nanoparticles (MNPs)
<b>Uniprot ID</b>	P30874
<b>Protein Families</b>	GPCR
<b>Protein Pathways</b>	Neuroactive ligand-receptor interaction
<b>Molecular Weight</b>	The human full length SSTR2 Protein has a MW of 41.2 kDa
<b>Delivery</b>	In Stock
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Sterility</b>	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 $\mu$ m) prior to use.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Somatostatin acts at many sites to inhibit the release of many hormones and other secretory proteins. The biologic effects of somatostatin are probably mediated by a family of G protein-coupled receptors that are expressed in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors having seven transmembrane segments and is expressed in highest levels in cerebrum and kidney.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated



### ELISA assay to evaluate SSTR2-MNPs

0.5 $\mu$ g HumanSSTR2-MNPs per well




Figure1. Elisa plates were pre-coated with 0.5 $\mu$ g/per well purified human SSTR2 full length membrane nanoparticles. Serial diluted anti-SSTR2 monoclonal antibody (BME100127) solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-SSTR2 monoclonal antibody binding with SSTR2 full length membrane nanoparticles is 86.2ng/ml.




Figure2. FACS analysis of SSTR2 MNPs

- Negative Control 1: SSTR2 full length membrane nanoparticles samples were stained only with Goat anti-human IgG 488 secondary antibody.
- Negative Control 2: Control membrane nanoparticles samples were stained with anti-SSTR2 antibody (BME100127) at 2 $\mu$ g/mL, followed by Goat anti-human IgG 488 secondary antibody.
- Negative Control 3: SSTR2 full length membrane nanoparticles samples were stained with anti-GPRC5D antibody (an irrelevant antibody) at 2 $\mu$ g/mL, followed by Goat anti-human IgG 488 secondary antibody.
- SSTR2 full length membrane nanoparticles samples were stained with anti-SSTR2 antibody (BME100127) at 2 $\mu$ g/mL, followed by Goat anti-human IgG 488 secondary antibody.

