

## **PRODUCT INFORMATION**

C-Flag&Strep Tag Tag

TAS2R38 **Target** 

**Synonyms** PTC; T2R38; T2R61; THIOT

Human TAS2R38-Strep full length protein-**Description** 

synthetic nanodisc

**Delivery** In Stock **Uniprot ID** P59533 **Expression Host HEK293** 

Storage & Shipping

**Background** 

**Protein Families** Druggable Genome, GPCR, Transmembrane

**Protein Pathways** Taste transduction

The human full length TAS2R38-Strep protein has **Molecular Weight** 

a MW of 37.9 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before Formulation & Reconstitution

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

A seven-transmembrane G protein-coupled receptor that controls the ability to taste glucosinolates, a family of bitter-tasting compounds found in plants of the Brassica sp Synthetic compounds phenylthiocarbamide (PTC) and 6-n-propylthiouracil (PROP) have been identified as ligands for this receptor and have been used to test the genetic diversity of this gene. Although several allelic forms of this gene

have been identified worldwide, there are two predominant common forms (taster and non-taster) found outside of Africa. These alleles differ at three nucleotide positions resulting in amino acid changes in the protein (A49P, A262V, and V296I) with the amino acid combination PAV identifying the taster variant (and AVI identifying

> Email: info@dimabio.com Website: www.dimabio.com

the non-taster variant).

Usage Research use only

Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)





## ELISA assay to evaluate TAS2R38-Strep-Nanodisc 0.2µg Human TAS2R38-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag TAS2R38-Strep-Nanodisc ( $0.2\mu g/per$  well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with TAS2R38-Strep-nanodisc is 8.159ng/ml.




Figure 2. Human TAS2R38-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

