

PRODUCT INFORMATION

| | |
|---|--|
| Target | TIMP2 |
| Synonyms | DDC8; CSC-21K |
| Description | Recombinant human TIMP2 Protein with C-terminal 6×His tag |
| Delivery | In Stock |
| Uniprot ID | P16035 |
| Expression Host | HEK293 |
| Tag | C-6×His tag |
| Molecular Characterization | TIMP2(Cys27-Pro220) 6×His tag |
| Molecular Weight | The protein has a predicted molecular mass of 22.6 kDa after removal of the signal peptide. |
| Purity | The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage&Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene is a member of the TIMP gene family. The proteins encoded by this gene family are natural inhibitors of the matrix metalloproteinases, a group of peptidases involved in degradation of the extracellular matrix. In addition to an inhibitory role against metalloproteinases, the encoded protein has a unique role among TIMP family members in its ability to directly suppress the proliferation of endothelial cells. As a result, the encoded protein may be critical to the maintenance of tissue homeostasis by suppressing the proliferation of quiescent tissues in response to angiogenic factors, and by inhibiting protease activity in tissues undergoing remodelling of the extracellular matrix. [provided by RefSeq, Jul 2008] |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1. Human TIMP2 Protein, His Tag on SDS-PAGE under reducing condition.

