Cat. No. PME100577

## **PRODUCT INFORMATION**

**Target** TLR3 TLR3 **Synonyms** 

Recombinant human TLR3 protein with C-terminal Description

6×His tag In Stock

**Delivery Uniprot ID** 015455 **Expression Host HEK293** 

Tag C-6×His Tag

Molecular

Storage & Shipping

**Background** 

TLR3(Ser24-Leu704) 6×His tag Characterization

The protein has a predicted molecular mass of **Molecular Weight** 

78.2 kDa after removal of the signal peptide. The apparent molecular mass of TLR3-His is

approximately 130-180 kDa due to glycosylation.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation &

- 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this gene is a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. They

recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity. The various TLRs exhibit different patterns of expression. This receptor is most abundantly expressed in placenta and pancreas, and is restricted to the dendritic subpopulation of

the leukocytes. It recognizes dsRNA associated with viral infection, and induces the activation of NF-kappaB and the production of type I interferons. It may thus play a role in host defense against viruses. Use of alternative polyadenylation sites to generate different length transcripts has been noted for this gene.

Email: info@dimabio.com Website: www.dimabio.com

[provided by RefSeq, Jul 2008]


Usage Research use only Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)





Figure 1. Human TLR3 Protein, His Tag on SDS-PAGE under reducing condition.



Email: info@dimabio.com Website: www.dimabio.com

SIMIABIO COLIFINITATION CONTRACTOR OF THE PROPERTY OF THE PROP

