

PRODUCT INFORMATION

Target	TNFRSF11A
Synonyms	Tumor necrosis factor receptor superfamily member 11A;ODFR;Receptor activator of NF-KB;CD265;RANK
Description	Recombinant human TNFRSF11A protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	Q9Y6Q6
Expression Host	HEK293
Tag	C-Human Fc Tag
Molecular Characterization	TNFRSF11A(Ile30-Pro212) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 46.2 kDa after removal of the signal peptide. The apparent molecular mass of TNFRSF11A-hFc is approximately 55-70 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptors can interact with various TRAF family proteins, through which this receptor induces the activation of NF-kappa B and MAPK8/JNK. This receptor and its ligand are important regulators of the interaction between T cells and dendritic cells. This receptor is also an essential mediator for osteoclast and lymph node development. Mutations at this locus have been associated with familial expansile osteolysis, autosomal recessive osteopetrosis, and Paget disease of bone. Alternatively spliced transcript variants have been described for this locus. [provided by RefSeq, Aug 2012]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human TNFRSF11A Protein, hFc Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

