

PRODUCT INFORMATION

| | |
|------------------------------|--|
| Target | TPO |
| Synonyms | Thrombopoietin;C-mpl ligand;Megakaryocyte colony-stimulating factor;Megakaryocyte growth and development factor;Myeloproliferative leukemia virus oncogene ligand;THPO |
| Description | Recombinant Human Thrombopoietin is produced by our Mammalian expression system and the target gene encoding Ser22-Gly353 is expressed with a 6His tag at the N-terminus, 6His tag at the C-terminus. |
| Delivery | In Stock |
| Uniprot ID | P40225 |
| Expression Host | HEK293 |
| Tag | N, C-6×His Tag |
| Molecular Characterization | Not available |
| Molecular Weight | 37.3 KDa |
| Purity | Greater than 95% as determined by reducing SDS-PAGE. |
| Formulation & Reconstitution | Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | Thrombopoietin (TPO) is a glycoprotein hormone which belongs to the EPO/TPO family. It produced by the liver and kidney which regulates the production of platelets. TPO stimulates the production and differentiation of megakaryocytes, the bone marrow cells that bud off large numbers of platelets. Lineage-specific cytokine affects the proliferation and maturation of megakaryocytes from their committed progenitor cells. It acts at a late stage of megakaryocyte development. It may be the major physiological regulator of circulating platelets. |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1. Greater than 95% as determined by reducing SDS-PAGE.

