

PRODUCT INFORMATION

|                              |   |
|------------------------------|---|
| Tag                          | C-Flag Tag  |
| Target                       | TRPA1   |
| Synonyms                     | ANKTM1; FEPS; FEPS1   |
| Description                  | Human TRPA1 full length protein-synthetic nanodisc  |
| Delivery                     | In Stock  |
| Uniprot ID                   | O75762  |
| Expression Host              | HEK293  |
| Protein Families             | Druggable Genome, Ion Channels: Transient receptor potential, Transmembrane   |
| Protein Pathways             | N/A   |
| Molecular Weight             | The human full length TRPA1 protein has a MW of 127.5 kDa   |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.  |
| Storage & Shipping           | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| Background                   | The structure of the protein is highly related to both the protein ankyrin and transmembrane proteins. This protein is activated by a large variety of structurally unrelated electrophilic and non-electrophilic chemical compounds. Electrophilic ligands activate TRPA1 by interacting with critical N-terminal Cys residues in a covalent manner, whereas mechanisms of non-electrophilic ligands are not well determined. May be a component for the mechanosensitive transduction channel of hair cells in inner ear, thereby participating in the perception of sounds. Probably operated by a phosphatidylinositol second messenger system. |
| Usage                        | Research use only   |
| Conjugate                    | Unconjugated  |






Figure1. Elisa plates were pre-coated with Flag Tag TRPA1-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with TRPA1-Nanodisc is 7.433ng/ml.



Figure2. Human TRPA1-Nanodisc, Flag Tag on SDS-PAGE

