

PRODUCT INFORMATION

|                              |   |
|------------------------------|---|
| Tag                          | C-Flag Tag  |
| Target                       | TRPC4   |
| Synonyms                     | HTRP-4, HTRP4, TRP4   |
| Description                  | Human TRPC4 full length protein-synthetic nanodisc  |
| Delivery                     | 6~8weeks  |
| Uniprot ID                   | Q9UBN4  |
| Expression Host              | HEK293  |
| Protein Families             | Ion Channels: Transient receptor potential  |
| Protein Pathways             | N/A   |
| Molecular Weight             | The human full length TRPC4 protein has a MW of 112.1kDa  |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.  |
| Storage&Shipping             | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| Background                   | This gene encodes a member of the canonical subfamily of transient receptor potential cation channels. The encoded protein forms a non-selective calcium-permeable cation channel that is activated by Gq-coupled receptors and tyrosine kinases, and plays a role in multiple processes including endothelial permeability, vasodilation, neurotransmitter release and cell proliferation. Single nucleotide polymorphisms in this gene may be associated with generalized epilepsy with photosensitivity. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Aug 2011] |
| Usage                        | Research use only   |
| Conjugate                    | Unconjugated  |

