Cat. No. PME100598



PRODUCT INFORMATION

UPA **Target**

Synonyms ATF;BDPLT5;QPD;u-PA;UPA;URK

Recombinant human UPA protein with C-terminal **Description**

6×His tag

Delivery In Stock **Uniprot ID** P00749 **HEK293 Expression Host** Tag C-6×His Tag

Molecular

Purity

Background

UPA(Ser21-Leu431) 6×His tag Characterization

The protein has a predicted molecular mass of

47.2 kDa after removal of the signal peptide. The apparent molecular mass of UPA-His is **Molecular Weight**

approximately 55-70 kDa due to glycosylation. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a secreted serine protease that converts plasminogen to plasmin. The encoded preproprotein is proteolytically processed to generate A and B polypeptide chains. These chains associate via a single disulfide bond to form the catalytically inactive high molecular weight urokinase-type plasminogen activator (HMW-uPA). HMW-uPA can

be further processed into the catalytically active low molecular weight urokinase-type plasminogen activator (LMW-uPA). This low molecular weight form does not bind to the urokinase-type

plasminogen activator receptor. Mutations in this gene may be associated with Quebec platelet disorder and late-onset Alzheimer's disease. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed. [provided by

> Email: info@dimabio.com Website: www.dimabio.com

RefSeq, Jan 2016]

Usage Research use only

Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)








Figure 1. Human UPA Protein, His Tag on SDS-PAGE under reducing condition.



