

PRODUCT INFORMATION

| | |
|------------------------------|---|
| Target | VIPR2 |
| Synonyms | VPAC2; VPAC2R; VIP-R-2; VPCAP2R; PACAP-R3; DUP7q36.3; PACAP-R-3; C16DUPq36.3 |
| Description | Recombinant human VIPR2 Protein with C-terminal human Fc tag |
| Delivery | In Stock |
| Uniprot ID | P41587 |
| Expression Host | HEK293 |
| Tag | C-Human Fc tag |
| Molecular Characterization | VIPR2(Glu24-Ile124) hFc(Glu99-Ala330) |
| Molecular Weight | The protein has a predicted molecular mass of 37.6 kDa after removal of the signal peptide. |
| Purity | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene encodes a receptor for vasoactive intestinal peptide, a small neuropeptide. Vasoactive intestinal peptide is involved in smooth muscle relaxation, exocrine and endocrine secretion, and water and ion flux in lung and intestinal epithelia. Its actions are effected through integral membrane receptors associated with a guanine nucleotide binding protein which activates adenylate cyclase. [provided by RefSeq, Aug 2011] |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1. Human VIPR2 Protein, hFc Tag on SDS-PAGE under reducing condition.

