

PRODUCT INFORMATION

Target	BTLA
Synonyms	BTLA;D272
Description	Recombinant mouse BTLA protein with C-terminal mFc tag
Delivery	In Stock
Uniprot ID	Q7TSA3
Expression Host	HEK293
Tag	C-Mouse Fc Tag
Molecular Characterization	Mouse BTLA(Glu30-Pro176) mFc(Pro99-Lys330)
Molecular Weight	The protein has a predicted molecular mass of 42.7 kDa after removal of the signal peptide. The apparent molecular mass of Mouse-BTLA-mFc is approximately 55-70 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Inhibitory receptor on lymphocytes that negatively regulates antigen receptor signaling via PTPN6/SHP-1 and PTPN11/SHP-2 (PubMed:12796776, PubMed:14652006). May interact in cis (on the same cell) or in trans (on other cells) with TNFRSF14 (PubMed:19915044). In cis interactions, appears to play an immune regulatory role inhibiting in trans interactions in naive T cells to maintain a resting state. In trans interactions, can predominate during adaptive immune response to provide survival signals to effector T cells (PubMed:19915044).
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Mouse BTLA Protein, mFc Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

