

## PRODUCT INFORMATION

|   |  |
|---|--|
| <b>Target</b>                           | NKG2D  |
| <b>Synonyms</b>                         | NK cell receptor D;CD314;Klrk1;Nkg2d   |
| <b>Description</b>                      | Recombinant mouse NKG2D protein with N-terminal human Fc tag   |
| <b>Delivery</b>                         | In Stock   |
| <b>Uniprot ID</b>                       | O54709   |
| <b>Expression Host</b>                  | HEK293   |
| <b>Tag</b>                              | N-Human Fc Tag   |
| <b>Molecular Characterization</b>       | hFc(Glu99-Ala330) Mouse NKG2D(Phe90-Val232)  |
| <b>Molecular Weight</b>                 | The protein has a predicted molecular mass of 42.6 kDa after removal of the signal peptide. The apparent molecular mass of hFc-mNKG2D is approximately 35-55 kDa due to glycosylation.   |
| <b>Purity</b>                           | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.   |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.  |
| <b>Storage&amp;Shipping</b>             | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.  |
| <b>Sterility</b>                        | Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.  |
| <b>Background</b>                       | Function as an activating and costimulatory receptor involved in immunosurveillance upon binding to various cellular stress-inducible ligands displayed at the surface of autologous tumor cells and virus-infected cells. Provides both stimulatory and costimulatory innate immune responses on activated killer (NK) cells, leading to cytotoxic activity. Acts as a costimulatory receptor for T-cell receptor (TCR) in CD8( ) T-cell-mediated adaptive immune responses by amplifying T-cell activation. Stimulates perforin-mediated elimination of ligand-expressing tumor cells. Signaling involves calcium influx, culminating in the expression of TNF-alpha. Participates in NK cell-mediated bone marrow graft rejection. May play a regulatory role in differentiation and survival of NK cells. Binds to ligands belonging to various subfamilies of MHC class I-related glycoproteins including RAET1A, RAET1B, RAET1C, RAET1D, RAET1E, H60 and MULT1.[UniProtKB/Swiss-Prot Function] |
| <b>Usage</b>                            | Research use only  |
| <b>Conjugate</b>                        | Unconjugated   |






Figure 1. Mouse NKG2D Protein, hFc Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

