

PRODUCT INFORMATION

TIM3 **Target**

HAVCR2;TIMD3;FLJ14428;KIM3 **Synonyms**

Recombinant mouse TIM3 protein with C-terminal **Description**

human Fc tag

Delivery In Stock **Uniprot ID** 08VIM0 **Expression Host HEK293**

Tag C-Human Fc Tag

Molecular

Background

Mouse TIM3(Arg20-Thr192) hFc(Glu99-Ala330) Characterization

The protein has a predicted molecular mass of 45.4 kDa after removal of the signal peptide. The **Molecular Weight**

apparent molecular mass of mTIM3-hFc is approximately 55-70 kDa due to glycosylation.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this gene belongs to the immunoglobulin superfamily, and TIM family of proteins. CD4-positive T helper lymphocytes can be divided into types 1 (Th1) and 2 (Th2) on the basis of their cytokine secretion patterns. Th1 cells are involved in cell-mediated immunity to intracellular pathogens and delayed-type


hypersensitivity reactions, whereas, Th2 cells are involved in the control of extracellular helminthic infections and the promotion of atopic and allergic diseases. This protein is a Th1-specific

cell surface protein that regulates macrophage activation, and inhibits Th1-mediated auto- and alloimmune responses, and promotes immunological tolerance. [provided by RefSeq,

> Email: info@dimabio.com Website: www.dimabio.com

Sep 2011]

Usage Research use only Conjugate Unconjugated








Figure 1. Mouse TIM3 Protein, hFc Tag on SDS-PAGE under reducing condition.

Email: info@dimabio.com Website: www.dimabio.com

