

**PRODUCT INFORMATION**

<b>Target</b>	TNFSF15
<b>Synonyms</b>	TI1;TI1a;Vegi;Tnlg1b
<b>Description</b>	Recombinant mouse TNFSF15 protein with N-terminal human Fc tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q5UBV8
<b>Expression Host</b>	HEK293
<b>Tag</b>	N-Human Fc Tag
<b>Molecular Characterization</b>	hFc(Glu99-Ala330) Mouse TNFSF15(Ala61-Leu252)
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 47.6 kDa after removal of the signal peptide. The apparent molecular mass of hFc-mTNFSF15 is approximately 55-70 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Sterility</b>	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
<b>Background</b>	Predicted to enable death receptor binding activity. Predicted to be involved in activation of cysteine-type endopeptidase activity involved in apoptotic process and positive regulation of cytokine production. Predicted to act upstream of or within activation of NF-kappaB-inducing kinase activity. Predicted to be located in plasma membrane. Used to study inflammatory bowel disease 16. Orthologous to human TNFSF15 (TNF superfamily member 15). [provided by Alliance of Genome Resources, Apr 2022]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated





Figure 1. Mouse TNFSF15 Protein, hFc Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

