

PRODUCT INFORMATION

Tag	N-Flag, C-Strep Tag
Target	N/A
Synonyms	N/A
Description	Negative control-synthetic nanodisc
Delivery	In Stock
Uniprot ID	N/A
Expression Host	HEK293
Protein Families	N/A
Protein Pathways	N/A
Molecular Weight	The protein has a MW of 42.8 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. Protect from light.
Formulation & Reconstitution	
Storage&Shipping	
Background	Negative control-synthetic nanodisc can be used as isotype control for nanodisc displaying membrane proteins in various applications. The protein in the nanodisc is a fusion protein, divided into three parts. N-terminal extracellular region is Flag tag. C-terminal intracellular region is EGFP-Strep tag. The transmembrane region of this protein utilizes the transmembrane region of platelet-derived growth factor receptor beta (PDGFRB).
Usage	Research use only
Conjugate	Fluorescent



ELISA assay to evaluate Negative control-Nanodisc
0.2 μ g Anti-Flag Antibody per well




Figure 1. Elisa plates were added with serial diluted N-Flag, C-Strep Tag Negative control-Nanodisc on an anti-Flag monoclonal antibody pre-coated (0.2 μ g/per well) plate. Strep tactin-HRP (1:5000) was used as detection antibody. From above data, the EC50 for Strep tactin-HRP binding with Negative control-Nanodisc is 283.0ng/ml.



Figure 2. Negative control-Nanodisc, N-Flag, C-Strep Tag on SDS-PAGE






Figure 3. WB analysis of Negative control-Nanodisc with Strep-Tactin HRP at 1/4000 dilution

