

**PRODUCT INFORMATION**

<b>Clone ID</b>	DM145
<b>Target</b>	CD46
<b>Synonyms</b>	CD46;AHUS2;MCP;MIC10;TLX;TRA2.10
<b>Host Species</b>	Rabbit
<b>Description</b>	PE-conjugated Anti-CD46 antibody(DM145); Rabbit mAb
<b>Delivery</b>	Under Development
<b>Uniprot ID</b>	P15529
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	Flow Cyt
<b>Recommended Dilutions</b>	Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Endotoxin</b>	Less than 1.0 EU/μg by the LAL method. For <1 EU/mg requirements, please contact us for customization.
<b>Formulation &amp; Reconstitution</b>	Liquid PBS with 0.05% Proclin300, 1% BSA
<b>Storage&amp;Shipping</b>	Store at 2°C-8°C for 6 months
<b>Sterility</b>	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 μm) prior to use. The protein encoded by this gene is a type I membrane protein and is a regulatory part of the complement system. The encoded protein has cofactor activity for inactivation of complement components C3b and C4b by serum factor I; which protects the host cell from damage by complement. In addition; the encoded protein can act as a receptor for the Edmonston strain of measles virus; human herpesvirus-6; and type IV pili of pathogenic Neisseria. Finally; the protein encoded by this gene may be involved in the fusion of the spermatozoa with the oocyte during fertilization. Mutations at this locus have been associated with susceptibility to hemolytic uremic syndrome. Alternatively spliced transcript variants encoding different isoforms have been described.
<b>Background</b>	
<b>Usage</b>	Research use only
<b>Conjugate</b>	PE-conjugated
<b>DIMA Disclaimer</b>	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr

