

## **PRODUCT INFORMATION**

Clone ID DM125 IL17RA **Target** 

**Synonyms** CD217;CDw217;IL-17RA;IL17R;CANDF5;hIL-17R

**Host Species** 

PE-conjugated Anti-IL17RA antibody(DM125); Description

Rabbit mAb

**Delivery** Under Development

**Uniprot ID** Q96F46 Rabbit IgG IgG type Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended Flow Cyt 1:100 **Dilutions** 

Formulation &

**DIMA Disclaimer** 

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Liquid PBS with 0.05% Proclin300, 1% BSA Reconstitution

Storage & Shipping Store at 2°C-8°C for 6 months

> Interleukin 17A (IL-17A) is a proinflammatory cytokine secreted by activated T-lymphocytes. It is a potent inducer of the maturation of CD34positive hematopoietic precursors into neutrophils. The transmembrane protein encoded by this gene (interleukin 17A receptor; IL17RA) is a ubiquitous type I membrane glycoprotein that binds with low affinity to interleukin 17A.

**Background** 

Interleukin 17A and its receptor play a pathogenic role in many inflammatory and autoimmune diseases such as rheumatoid arthritis. Like other cytokine receptors; this receptor likely has a multimeric structure. Alternative splicing results in multiple transcript variants encoding different

isoforms.

**Usage** Research use only

Conjugate PE-conjugated

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are

actively scrutinizing all patent application to

ensure no IP infringement.





