

PRODUCT INFORMATION

Target	S protein RBD
Synonyms	S protein RBD;Spike glycoprotein Receptor-binding domain;S glycoprotein RBD;Spike protein RBD;COVID-19
Description	Recombinant SARS-CoV-2 (2019-nCoV) S protein RBD with C-terminal 6×His tag
Delivery	In Stock
Uniprot ID	P0DTC2
Expression Host	HEK293
Tag	C-6×His Tag
Molecular Characterization	S protein RBD(Arg319-Phe541) 6×His tag
Molecular Weight	The protein has a predicted molecular mass of 37-38 kDa after removal of the signal peptide.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) also known as Covid19 (2019 Novel Coronavirus) is a virus that causes illnesses ranging from the common cold to severe diseases. The spike protein is a type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which accounts for recognizing the cell surface receptor, ACE2. S2 contains basic elements needed for the membrane fusion. Recent publications indicate that S1-RBD domain can induce virus neutralizing-antibody and T cell response.
Usage	Research use only
Conjugate	Unconjugated






Figure 1. SARS-CoV-2 (2019-nCoV) S protein RBD, His Tag on SDS-PAGE under reducing condition.

S-RBD, His Tagged protein ELISA

0.2 µg S-RBD, His Tagged protein per well




Figure 2. ELISA plate pre-coated by 2 µg/ml (100 µl/well) SARS-CoV-2 (2019-nCoV) S protein RBD, His tagged protein (PME100036) can bind its native ligand Human ACE2, hFc tagged protein PME100073 in a linear range of 1.96-31.72 ng/ml.

